

GLP-1/DIABETIC MEDICATION GUIDELINES

GLP-1 medications: both injections and pill form***TAKEN FOR DIABETES AND/OR WEIGHT LOSS***

For patients scheduled at **the ASC** medication must be held for **at least 1 week**

For patients scheduled at the **Corewell Hospital** medication must be held **2 weeks**

- **Ozempic** (shot)
- **Wegovy** (shot)
- **Trulicity** (shot)
- **Victoza** (shot)
- **Saxenda** (shot) ****TAKEN FOR DIABETES AND/OR WEIGHT LOSS****
- **Byetta** (shot)
- **Bydureon BCise** (shot)
- **Mounjaro** (shot)
- **Zepbound** (shot)
- **Adlyxin** (shot)
- **Rybelsus** (pill)
- **Adipex (phentermine)** ****NOT A DIABETIC MED (used for appetite suppression/weight loss)****

Diabetic medications, which are **not GLP-1** and should be stopped according to the diabetic guidelines:

For patients scheduled at **the ASC** and **Corewell Hospital** medication **hold 3 days prior**

- **Invokana**
- **Farxiga**
- **Jardiance**

Oral meds: ½ morning dose and hold evening dose day before, do not take day of procedure

- | | |
|---|--------------------|
| • Metformin (can also be taken for PCOS) | • Starlix |
| • Welchol | • Prandin |
| • Cycloset | • Amaryl |
| • Januvia | • Glucotrol |
| • Onglyza | • Glyburide |
| • Tradjenta | • Actos |
| • Nesina | • Avandia |

Insulin injections: ½ morning dose and ½ evening dose *day before*

Long-acting insulin (½ dose *morning of*): Lantus, Basaglar, Levemir, Toujeo, Tresiba

Short-acting insulin (STOP *day of* unless continuous pump): Novolog, Novolin, Humalog, Apidra