

GLP/DIABETIC MEDICATION GUIDELINES

Center for Digestive Health - www.Center4DigestiveHealth.com - (248) 844-9710

Hold 7 Days Before Procedure	Hold 3 Days Before Procedure	1 Day before: ½ morning dose and hold evening dose, DO NOT take day of procedure
Any GLP on Clinical Trial	Canagliflozin (Invokana)	Acarbose (Precose)
Phentermine (Adipex-P)	Dapagliflozin (Farxiga)	Alogliptin (Nesina)
Naltrexone/Bupropion (Contrave)	Empagliflozin/Metformin (Synjardy)	Bromocriptine (Cycloset)
Dulaglutide (Trulicity)	Empagliflozin (Jardiance)	Colesevelam (Welchol)
Exenatide (Byetta, Bydureon)	Empagliflozin/Linagliptin/Metformin (Trijardy)	Glimepiride (Amaryl)
Liraglutide (Saxenda, Victoza)	Oral daily Orforglipron (Foundayo)	Glipizide (Glucotrol)
Lixisenatide (Adlyxin)	Oral daily Semaglutide (Rybelsus, Wegovy)	Glyburide (Glynase)
Semaglutide Injection (Ozempic, Wegovy)		Linagliptin (Tradjenta)
Tirzepatide (Mounjaro, Zepbound)		Metformin (Fortemet, Glucophage, Glumetza)
		Nateglinide (Starlix)
		Pioglitazone (Actos)
		Repaglinide (Prandin)
		Rosiglitazone (Avandia)
		Saxagliptin (Onglyza)
		Sitagliptin (Januvia)
		Sitagliptin/Metformin (Janumet)

(248) 844-9710

Center4DigestiveHealth.com

INSULINS		FAILURE TO HOLD THESE MEDICATIONS WILL RESULT IN CANCELLATION OF YOUR PROCEDURE DUE TO ANESTHESIA RISKS
Long Acting- Administer 1/2 of normal dose the morning of the procedure	Short/Intermediate Acting- DO NOT administer the day of (unless on a continuous pump)	
Degludec (Tresiba)	Aspart (Novolog)	
Detemir (Levemir)	Glulisine (Apidra)	
Glargine (Basaglar, Lantus, Rezvoglar, Semglee, Toujeo)	Lispro (Humalog)	
	NPH (Humulin, Novolin)	
	Regular (Humulin, Novolin)	

updated 7/17/2026